

This IssueViews **35,614** | Citations **0** | Altmetric **1239**

PDF

 Full Text Share

Original Investigation

March 24/31, 2020


Association of Daily Step Count and Step Intensity With Mortality Among US Adults

Pedro F. Saint-Maurice, PhD¹; Richard P. Troiano, PhD²; David R. Bassett Jr, PhD³; [et al](#)

Author Affiliations

¹Division of Cancer Epidemiology and Genetics, National Cancer Institute, Rockville, Maryland²Division of Cancer Control and Population Sciences, National Cancer Institute, Rockville, Maryland³Department of Kinesiology, Recreation, and Sport Studies, University of Tennessee, Knoxville⁴Division of Nutrition, Physical Activity, and Obesity, National Center for Chronic Disease Prevention and Health Promotion, Centers for Disease Control and Prevention, Atlanta, Georgia⁵Epidemiology and Population Science Laboratory, National Institute on Aging, Bethesda, Maryland

JAMA. 2020;323(12):1151-1160. doi:10.1001/jama.2020.1382

 Related Articles

Key Points

Question What are the associations between daily step counts and step intensity with mortality among US adults?

Findings In this observational study that included 4840 participants, a greater number of steps per day was significantly associated with lower all-cause mortality (adjusted hazard ratio for 8000 steps/d vs 4000 steps/d, 0.49). There was no significant association between step intensity and all-cause mortality after adjusting for the total number of steps per day.

Our website uses cookies to enhance your experience. By continuing to use our site, or clicking "Continue," you are agreeing to our [Cookie Policy](#) | [Continue](#)

Abstract

Importance It is unclear whether the number of steps per day and the intensity of stepping are associated with lower mortality.

Objective Describe the dose-response relationship between step count and intensity and mortality.

Design, Setting, and Participants Representative sample of US adults aged at least 40 years in the National Health and Nutrition Examination Survey who wore an accelerometer for up to 7 days (from 2003-2006). Mortality was ascertained through December 2015.

Exposures Accelerometer-measured number of steps per day and 3 step intensity measures (extended bout cadence, peak 30-minute cadence, and peak 1-minute cadence [steps/min]). Accelerometer data were based on measurements obtained during a 7-day period at baseline.

Main Outcomes and Measures The primary outcome was all-cause mortality. Secondary outcomes were cardiovascular disease (CVD) and cancer mortality. Hazard ratios (HRs), mortality rates, and 95% CIs were estimated using cubic splines and quartile classifications adjusting for age; sex; race/ethnicity; education; diet; smoking status; body mass index; self-reported health; mobility limitations; and diagnoses of diabetes, stroke, heart disease, heart failure, cancer, chronic bronchitis, and emphysema.

Results A total of 4840 participants (mean age, 56.8 years; 2435 [54%] women; 1732 [36%] individuals with obesity) wore accelerometers for a mean of 5.7 days for a mean of 14.4 hours per day. The mean number of steps per day was 9124. There were 1165 deaths over a mean 10.1 years of follow-up, including 406 CVD and 283 cancer deaths. The unadjusted incidence density for all-cause mortality was 76.7 per 1000 person-years (419 deaths) for the 655 individuals who took less than 4000 steps per day; 21.4 per 1000 person-years (488 deaths) for the 1727 individuals who took 4000 to 7999 steps per day; 6.9 per 1000 person-years (176 deaths) for the 1539 individuals who took 8000 to 11999 steps per day; and 4.8 per 1000 person-years (82 deaths) for the 919 individuals who took at least 12000 steps per day. Compared with taking 4000 steps per day, taking 8000 steps per day was associated with significantly lower all-cause mortality (HR, 0.49 [95% CI, 0.44-0.55]), as was taking 12000 steps per day (HR, 0.35 [95% CI, 0.28-0.45]). Unadjusted incidence density for all-cause mortality by peak 30 cadence was 32.9 per 1000 person-years (406 deaths) for the 1080 individuals who took 18.5 to 56.0 steps per minute; 12.6 per 1000 person-years (207 deaths) for the 1153 individuals who took 56.1 to 69.2 steps per minute; 6.8 per 1000 person-years (124 deaths) for the 1074 individuals who took 69.3 to 82.8 steps per minute; and 5.3 per 1000 person-years (108 deaths) for the 1037 individuals who took 82.9 to 149.5 steps per minute. Greater step intensity was not significantly associated with lower mortality after adjustment for total steps per day (eg, highest vs lowest quartile of peak 30 cadence: HR, 0.90 [95% CI, 0.65-1.27]; *P* value for

Our website uses cookies to enhance your experience. By continuing to use our site, or clicking "Continue," you are agreeing to our [Cookie Policy](#) | [Continue](#)

Conclusions and Relevance Based on a representative sample of US adults, a greater number of daily steps was significantly associated with lower all-cause mortality. There was no significant association between step intensity and mortality after adjusting for total steps per day.



Full Text

Read More About

Cardiology

Physical Activity

Lifestyle Behaviors



Coronavirus Resource Center

SCHEDULED MAINTENANCE

Our websites may be periodically unavailable between 11:00PM CT April 18, 2020 and 4:00PM CT April 19, 2020 for regularly scheduled maintenance.

Trending

Opinion

Return of Research Results to Study Participants

August 7, 2018

Opinion

Appropriate Antibiotic Prescribing

May 3, 2016

Research

Our website uses cookies to enhance your experience. By continuing to use our site, or clicking "Continue," you are agreeing to our [Cookie Policy](#) | [Continue](#)

March 27, 2020

Select Your Interests

JOB LISTINGS ON JAMA CAREER CENTER®

Deputy Editor for Medicine

Worldwide (Remote)

Pulmonologist- Physician- Sioux City

Sioux City, Iowa

Pulmonologist

Buckhannon, West Virginia

California Pulmonary Critical Care Job

Carmel-by-the-Sea, California

Locums Job in CA for Critical Care Physician

Moreno Valley, California

See more at JAMA Career Center

Trending

Long-term Sequelae of Nonobstructive Chronic Bronchitis

JAMA Internal Medicine | *Opinion* | March 2, 2020

Nonobstructive Chronic Bronchitis and Respiratory Health Outcomes in Adults

JAMA Internal Medicine | *Research* | March 2, 2020

Association of Step Volume and Intensity With All-Cause Mortality in Older Women

JAMA Internal Medicine | *Research* | August 1, 2019

Our website uses cookies to enhance your experience. By continuing to use our site, or clicking "Continue," you are agreeing to our [Cookie Policy](#) | [Continue](#)